PoultryStar® me

Healthy gut – strong bird!



PoultryStar® me is a well-defined, poultry-specific, multi-genus synbiotic product that promotes beneficial gut microbiota through the combined action of specially selected probiotic microorganisms and prebiotic fructooligosaccharides.

PoultryStar® me is designed to improve gut health of poultry and to make day-old chick and birds of all ages less susceptible to be colonised by pathogens leading to improved animal health and food safety.

Mode of action

- **Our patented probiotic strains** were thoroughly investigated and selected due to their superior probiotic features and synergistic effects. They demonstrated the best efficacy in the competitive exclusion of pathogenic bacteria. The probiotic strains were isolated out of the gut of healthy birds to increase the probiotic-host compatibility.
- **Prebiotic fructooligosaccharides** (FOS) serve as a nutritional source specific for beneficial bacteria from the genera *Lactobacillus* and *Bifidobacterium*.

Main benefits

- Fast establishment of beneficial gut microbiota
- Re-establishment of balanced gut microbiota post antibiotic application
- Reduction of enteric opportunistic and pathogenic bacteria
- Reduces incidence of wet litter and pasty vent
- No negative side effects, no withdrawal period

PoultryStar® me is recommended to be used in the following conditions:

- Bacterial challenge
- Necrotic enteritis
- Wet droppings
- Pasty vent
- Vaccination
- Feed changes
- Extreme temperature
- High humidity
- High stocking densities
- Transportation

Learn more at
dsm-firmenich.com/anh

dsm-firmenich

PoultryStar® me

Healthy gut – strong bird!

Application

PoultryStar® me is mixed directly into finished feed.

PoultryStar® me can be pelleted up to temperatures of 85 °C.

The use of PoultryStar® me is especially beneficial to (re)-establish, and develop the microbiota in poultry post-hatching, during phases of intestinal stress and periods of high pathogenic challenge. Stress influences the gut microbiota negatively leading to dysbiosis and reduced performance.

Animal categories and species: chickens for fattening, chickens reared for laying (including laying period excl. EU), minor poultry species.

Dosage

0.5 – 1 kg/ ton of feed

Packing

PoultryStar® me is available in 20 kg bags and 25 kg corrugated cartons with polyethylene inlet.

Stability and storage

PoultryStar® me must be stored in a cool and dry place in its original packaging and away from direct sunlight. The minimum shelf life is 12 months from the production date if stored in unopened bags and according to the recommendations of the manufacturer.

Disclaimer: dsm-firmenich has used diligent care to ensure that the information provided herein is accurate and up-to-date, however, dsm-firmenich makes no representation or warranty, either expressly or implied, of the accuracy, reliability, or completeness thereof. The information provided herein contains scientific and product information for business to business use and does not constitute or provide scientific or medical advice, diagnosis, or recommendation for treatment. Country or region-specific information should be considered when labeling or advertising to the final consumer. In no event shall dsm-firmenich be liable for any damages arising from or reliance upon, or use of, any information provided herein. The content of this document is subject to change without further notice. Please contact your local dsm-firmenich representative for further details. All trademarks listed in this document are either (registered) trademarks of, or trademarks licensed by, the dsm-firmenich group of companies in the Netherlands and/or other countries, unless explicitly stated otherwise.

© dsm-firmenich, 2024.

PI_PoultryStarMe_EN_0224_ESL