
Effects of mycotoxins

T-2, DON, Ergots
- Decreased feed intake
- Dermal and oral lesions
- Feed refusal
- Vomiting
- Impaired growth

T-2, DON, AFB_1, OTA, FUM, Endotoxins
- Intestinal hemorrhages
- Kidney damage
- Pale and fatty liver
- Porcine pulmonary edema (PPE)
- Increased water consumption
- Fever – endotoxin-associated diseases

ZEN, T-2, DON, Ergots
- Irregular heats
- Abortion
- Pseudo pregnancy
- Low conception rates
- Ovarian cysts
- Embryonic Loss
- Tail necrosis
- Nymphomania
- Hypertrophy of the uterus
- Shrunken udder/ Agalactia
- Stillbirths



AFB_1, T-2, OTA
- Diarrhea
- Blood in faeces and urine
- Inflammation of bladder and kidneys

T-2, Ergots
- Dermal and oral lesions
- Vasoconstriction (necrosis)

T-2, DON, AFB_1, OTA, FUM
- Decreased performance
- Immunosuppression
- Pancreatic necroses

DON: Deoxynivalenol
ZEN: Zearalenone
AFB_1: Aflatoxin B_1
FUM: Fumonisins
T-2: T-2 Toxin
HT-2: HT-2 Toxin
NIV: Nivalenol
DAS: Diacetoxyscirpenol
OTA: Ochratoxin A
Ergots: Ergot Alkaloids

Learn more at
dsm-firmenich.com/anh

dsm-firmenich

PO_MTX_S_EN_1223_A2_KNA ©Copyright. dsm-firmenich 2024